



RESEARCH ARTICLE

10.1029/2019GC008809

Gravity, Topography, and Melt Generation Rates From Simple 3-D Models of Mantle Convection

Matthew E. Lees^{1,2} , John F. Rudge¹ , and Dan McKenzie¹

¹Bullard Laboratories, Department of Earth Sciences, University of Cambridge, Cambridge, UK, ²Now at School of Earth, Energy, and Environmental Sciences, Stanford University, Stanford, CA, USA

Key Points:

- Convection at high Rayleigh numbers has a spoke pattern planform with hot rising sheets (spokes) connected to upwelling plumes (hubs)
- Surface gravity and topography are most sensitive to the hubs of the pattern, but melt generation has additional sensitivity to the spokes
- Patterns of volcanism across North Africa and the Middle East are suggestive of spoke pattern convection beneath the plate

Supporting Information:

- Supporting Information S1

Correspondence to:

J. F. Rudge,
jfr23@cam.ac.uk

Citation:

Lees, M. E., Rudge, J. F., & McKenzie, D. (2020). Gravity, topography, and melt generation rates from simple 3-D models of mantle convection. *Geochemistry, Geophysics, Geosystems*, 21, e2019GC008809. <https://doi.org/10.1029/2019GC008809>

Received 7 NOV 2019

Accepted 6 MAR 2020

Accepted article online 10 MAR 2020

Abstract Convection in fluid layers at high Rayleigh number ($Ra \sim 10^6$) have a spoke pattern planform. Instabilities in the bottom thermal boundary layer develop into hot rising sheets of fluid, with a component of radial flow toward a central upwelling plume. The sheets form the “spokes” of the pattern, and the plumes the “hubs.” Such a pattern of flow is expected to occur beneath plate interiors on Earth, but it remains a challenge to use observations to place constraints on the convective planform of the mantle. Here we present predictions of key surface observables (gravity, topography, and rates of melt generation) from simple 3-D numerical models of convection in a fluid layer. These models demonstrate that gravity and topography have only limited sensitivity to the spokes and mostly reflect the hubs (the rising and sinking plumes). By contrast, patterns of melt generation are more sensitive to short-wavelength features in the flow. There is the potential to have melt generation along the spokes but at a rate which is relatively small compared with that at the hubs. Such melting of spokes can only occur when the lithosphere is sufficiently thin ($\lesssim 80$ km) and mantle water contents are sufficiently high ($\gtrsim 100$ ppm). The distribution of volcanism across the Middle East, Arabia, and Africa north of the equator suggests that it results from such spoke pattern convection.

1. Introduction

What is the planform of mantle convection? The largest, and most obvious, planform in the convection system is that associated with plate motions, which involves horizontal scales as large as 10,000 km (e.g., the Pacific plate). However, it is also clear that convection takes place at shorter horizontal scales. This scale of convection manifests in hot spot volcanism and the swells and troughs in gravity and topography observed at wavelengths of around 1,000 to 2,000 km (Crosby et al., 2006; Crosby & McKenzie, 2009; McKenzie, 1994). A well-cited example of this comes from Africa (Burke, 1996; Holmes, 1965; Jones et al., 2012; McKenzie & Weiss, 1975), which shows a clear pattern of swells and troughs across the continent. The strong correlation between the patterns of gravity and topography, and in particular the characteristic ratio of around 50 mGal km⁻¹ between the two, has been used to strongly argue for convective support of the topography (Crosby et al., 2006; Jones et al., 2012; McKenzie, 1994).

In theory, maps of gravity and topography should provide information on the planform of mantle convection. But extracting that information is challenging. While gravity provides information on density variations within the Earth, the process of inverting gravity data for density is highly nonunique. However, density variations within the Earth on length scales of hundreds of kilometers are not arbitrary but are controlled by the fluid dynamics of convection.

The aim of the present manuscript is to get a better understanding of the surface expressions of mantle convection from a series of the simplest possible numerical simulations of convection and to compare these with geophysical and geological observations. The fluid dynamical problem is one that has been extensively studied and discussed in the literature: Rayleigh-Bénard thermal convection in a 3-D rectangular box with a fluid of constant viscosity at Rayleigh numbers around 10^6 . This problem was first studied using laboratory experiments in large aspect ratio tanks, using a layer of silicone oil whose depth was a few centimeters (Busse & Whitehead, 1971, 1974; Richter & Parsons, 1975). These experiments showed that the planform of the convection was a spoke pattern, with hot rising plumes joined to each other by hot sheets near the lower thermal boundary layer and cold sinking plumes by cold sheets near the top thermal boundary layer. White (1988) carried out similar experiments using a fluid whose viscosity was a strong function of temperature and showed that at high Rayleigh number the convective planform was also a spoke pattern.

©2020. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

It is now possible to carry out three-dimensional time-dependent fully resolved numerical experiments at Rayleigh numbers of 10^5 – 10^7 , and the results of many such experiments have been reported (Christensen & Harder, 1991; Galsa & Lenkey, 2007; Houseman, 1990; Larsen et al., 1997; Tackley, 1993, 1996; Sotin & Labrosse, 1999; Zhong, 2005; Vilella & Deschamps, 2017, 2018; Weinstein & Christensen, 1991). Such experiments have an important advantage over tank experiments, because it is straightforward to calculate the geophysical observables from the numerical solutions.

We carried out our calculations with an infinite Prandtl number fluid and used the Boussinesq approximation throughout. We used large aspect ratio boxes, similar to those previously used in tank experiments, to allow the convective circulation to determine its own planform rather than being dominated by the lateral boundaries. We use our experiments to show which features of the observations are readily explained by the simplest models and which features are not. We do not attempt to construct a realistic model of the Earth. In this respect our aim is the same as that of the early tank experiments and, in addition, to extend them to encompass the observables: the gravity field, surface deformation, and melt generation. In particular, by allowing the density variations to arise naturally from fluid dynamics, such experiments allow an exploration of short-wavelength (< 100 km) temperature variations, which are likely to be most clearly expressed by volcanism.

Two effects that are known to be important in the Earth are not taken into account in the simple constant viscosity Boussinesq model we use. The first is the variation of viscosity with temperature. It is this effect that produces plates, and therefore our modeling does not include the dynamics of plate motions. The other effect is viscous dissipation, which is intimately related to vertical density variations that result from lithostatic pressure (Jarvis & McKenzie, 1980; Schubert et al., 2001; Spiegel & Veronis, 1960). Viscous heat generation has little effect on the circulation even when the relevant term in the equations cannot be neglected. Such heating occurs in boundary layers where temperature gradients are large. As a result the entropy and potential temperature (section A5) is little affected (Jarvis & McKenzie, 1980), though the convection becomes more time dependent.

The approach taken here complements a popular alternative approach to modeling gravity and topography using information from seismic tomography (Flament et al., 2013; Hager & Richards, 1989). In such studies, estimates of density variations within the mantle are inferred from tomography and used to make predictions of gravity and dynamic topography. Though these studies have had some success at predicting the very long wavelength ($> 6,000$ km) features of Earth's gravity, they depend on knowing the relationship between density ρ and the seismic velocities V_p and V_s , and also on a rheological model of the mantle. Furthermore, since these calculations are based on seismic tomography, they are limited by its resolution, which is not yet sufficient to map rising and sinking plumes in the upper mantle. Our approach also departs from the common assumption of many Earth scientists who believe the convective planform of mantle convection consists solely of plumes and the plates. This assumption arises from the work of Wilson (1963) and Morgan (1971), who showed that the relative motion between major volcanic centers beneath plate interiors was sufficiently slow that they could be used to define a single worldwide reference frame. Their ideas have been enormously influential. But they are based on an intuitive conception about the planform of high Rayleigh number convection, rather than on fluid dynamical experiments. They also predate our understanding of polybaric melt generation.

The manuscript is organized as follows. Section 2 describes the fluid dynamical simulations, and how they are scaled to parameter values appropriate for the Earth's mantle. Section 3 discusses the predicted gravity, topography, and their spectral properties. Section 4 discusses melt generation. Section 5 compares the results from the fluid dynamical experiments with the observed gravity and topography and with the volcanism of Africa and the Middle East, and conclusions follow in section 6. An appendix provides further technical details on the simulations and data processing.

2. Numerical Experiments

We ran 12 numerical experiments of isoviscous thermal convection in a rectangular box. Temperature was fixed at the top and bottom boundaries. To examine the influence of dynamical boundary conditions, runs were made for all combinations of freely slipping or rigid boundary conditions on the top and bottom boundaries. Reflection boundary conditions were applied at the side boundaries.




Figure 1. Artificial shadowgraphs of the 12 numerical experiments of thermal convection in an $8 \times 8 \times 1$ rectangular box. Each column is at a given Rayleigh number, and each row is at a given choice of boundary conditions for the Stokes flow. In each case the first word corresponds to the top boundary, and the second word to the bottom boundary; so free-rigid refers to a free-slip top and rigid bottom boundary condition.

All convection simulations were performed using v2.01 of the ASPECT mantle convection code (Bangerth et al., 2018; Dannberg & Heister, 2016; Heister et al., 2017). The code was used to solve the dimensionless versions of the Boussinesq governing equations of thermal convection in an $8 \times 8 \times 1$ rectangular box, through a small modification of the “convection-box” example discussed in the ASPECT manual. The governing equations in dimensional form are

$$\nabla \cdot \mathbf{v} = 0, \quad (1)$$

$$-\nabla P + \eta \nabla^2 \mathbf{v} = -\rho_0 g \alpha \theta \hat{\mathbf{z}}, \quad (2)$$

$$\frac{D\theta}{Dt} = \kappa \nabla^2 \theta, \quad (3)$$

where \mathbf{v} is the velocity, P is the difference in pressure from hydrostatic, θ is potential temperature, ρ_0 is the reference mantle density, α is the thermal expansivity, η is the viscosity, and κ is the thermal diffusivity.

Table 1
Common Parameter Values for All Runs

Quantity	Symbol	Value
Thermal expansivity	α	$4.0 \times 10^{-5} \text{ K}^{-1}$
Acceleration due to gravity	g	9.81 m s^{-2}
Layer thickness	d	600 km
Reference mantle density	ρ_0	$3,300 \text{ kg m}^{-3}$
Thermal conductivity	k	$3.7 \text{ W m}^{-1} \text{ K}^{-1}$
Specific heat capacity	C_p	$1.3 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

Note. These values are appropriate for the top ~ 100 km of the upper mantle beneath the lithosphere.

We assume constant thermal conductivity, constant heat capacity, constant viscosity, and constant thermal expansivity. In dimensionless form the governing equations are

$$\nabla \cdot \mathbf{v} = 0, \quad (4)$$

$$-\nabla P + \nabla^2 \mathbf{v} = -\text{Ra} \theta \hat{\mathbf{z}}, \quad (5)$$

$$\frac{D\theta}{Dt} = \nabla^2 \theta, \quad (6)$$

where all lengths have been scaled by the layer depth d , time by the diffusion time d^2/κ , pressure by $\eta\kappa/d^2$, and potential temperature by the potential temperature difference ΔT_p across the layer. Just one dimensionless parameter describes this simple system, the Rayleigh number, defined by

$$\text{Ra} = \frac{\rho_0 g \alpha \Delta T_p d^3}{\eta \kappa}. \quad (7)$$

Runs were performed at three different Rayleigh numbers: $\text{Ra} = 10^5$, 3×10^5 , and 10^6 . A uniform resolution of 32 cells in the vertical and 256 cells in the horizontal was specified for all simulations. Quadratic finite elements were used for temperature and velocity, and linear finite elements for pressure. Simulations were

Table 2
Parameter Values and Magnitudes of Observables for Each Simulation Which Yields a Lithospheric Thickness of 100 km and an Interior Potential Temperature of 1315 °C

BC	Ra	η (Pa s)	MBL (km)	ΔT_p (K)	h_{RMS} (m)	Δg_{RMS} (mGal)
Rigid-rigid	10^6	2.8×10^{20}	66.8	855	307	16.3
Rigid-rigid	3×10^5	1.3×10^{21}	53.8	1,188	636	32.9
Rigid-rigid	10^5	5.3×10^{21}	37.2	1,648	1,300	63.7
Free-rigid	10^6	2.3×10^{20}	80.1	709	244	11.2
Free-rigid	3×10^5	1.1×10^{21}	71.2	1,002	495	21.0
Free-rigid	10^5	4.4×10^{21}	59.4	1,349	958	37.1
Rigid-free	10^6	2.1×10^{20}	66.7	660	325	15.5
Rigid-free	3×10^5	1.0×10^{21}	54.6	935	672	32.6
Rigid-free	10^5	4.2×10^{21}	38.6	1,307	1,333	65.2
Free-free	10^6	1.4×10^{20}	82.8	442	235	10.1
Free-free	3×10^5	7.0×10^{20}	74.7	651	476	19.8
Free-free	10^5	3.0×10^{21}	64.6	912	943	37.7

Note. Columns from left to right: BC, boundary conditions at top-bottom; Ra, the Rayleigh number; η , viscosity; MBL, thickness of the mechanical boundary layer; ΔT_p , potential temperature difference across the fluid layer; h_{RMS} , root-mean-square of the dynamic topography at the top of the convecting region (as plotted in Figure 4); Δg_{RMS} , root-mean-square of the gravity anomaly at the top of the convecting region (as plotted in Figure 3).




Figure 2. Profiles of horizontally averaged potential temperature. Zero depth represents the top of the convecting region (the base of the mechanical boundary layer). The four panels show the four choices of boundary condition. The dimensional scaling is such that the interior potential temperature is 1315 °C and the lithospheric thickness is 100 km.

run until the system reached a quasi steady state, which was monitored by examining the behavior of the mean temperature and RMS velocity over time. Each simulation ran for a minimum of 6 times the thermal time constant for the layer ($= 6 d^2 / (\pi^2 \kappa)$).

Snapshots of the 12 experiments are shown in Figure 1. The images are made to mimic the shadowgraph visualization technique commonly used in laboratory experiments (Busse & Whitehead, 1971, 1974; Richter & Parsons, 1975; White, 1988; Whitehead & Parsons, 1977). To make a shadowgraph in the laboratory, light is shone from below through the layer of fluid and projected onto a screen. Refraction causes the light to focus and defocus according to the temperature variations within the fluid through which it passes. The effect is to make upwellings appear dark and downwellings appear bright. Mathematically, a shadowgraph produces a plot of the Laplacian of the vertically averaged temperature (Jenkins, 1988; Travis et al., 1990), and this is how the images in Figure 1 were computed.

The planform of convection at these Rayleigh numbers is spoke pattern (Busse & Whitehead, 1974). Hot and cold plumes develop from the top and bottom of the layer: the “hubs” of the planform. Hot and cold sheets, the “spokes” of the planform, radiate from the plumes. These sheets are formed by instabilities in the top and bottom boundary layers, and the shearing associated with the plumes suppresses instabilities with other geometries. The planform is time dependent.

Several well-known features stand out from the shadowgraph pictures. The first is that as the Rayleigh number increases the thicknesses of the upwellings and downwellings become narrower. For the free-free




Figure 3. Gravity anomalies at the top of the convecting box for the $Ra = 10^6$ simulations. The region shown is approximately the same size as the region outlined by the thick black line in Figure 11. Each panel shows a different combination of boundary conditions. In each case the first word corresponds to the top boundary, and the second word to the bottom boundary; so free-rigid refers to a free-slip top and rigid bottom boundary condition.

and rigid-rigid cases the upwellings and downwellings are symmetric, but there is a notable asymmetry in the nature of upwellings and downwellings in the free-rigid and rigid-free cases (Kvernfold, 1979; Weinstein & Christensen, 1991). Also notable is the change in horizontal distance between upwelling and downwelling with boundary condition: This distance is shorter for rigid-rigid simulations than for free-free simulations.

Our main interest here is the surface observables associated with mantle convection. The simulations were performed using dimensionless variables, with the dimensionless Rayleigh number as the only control parameter. To make predictions about observable quantities, these simulations must be scaled appropriately to Earth-like values. For most quantities we can simply use typical mantle estimates, and these values




Figure 4. Dynamic topography at the top of the convecting box for the $Ra = 10^6$ simulations.

are given in Table 1. We consider here just upper mantle convection and so choose as an appropriate layer thickness $d = 600$ km to scale all lengths.

Choosing an appropriate scaling of temperature is less straightforward. The numerical simulations essentially provide dimensionless potential temperatures. The potential temperature is the temperature that the mantle material would have if it were moved to the Earth's surface isentropically and without melting (McKenzie, 1970; section A5).

The scaling of the dimensionless temperature is required to satisfy two conditions: The first is that the average interior temperature must correspond to a mantle potential temperature of 1315°C . This choice ensures that the thickness of the oceanic crust is 7 km, generated by isentropic decompression beneath a spreading ridge using the parametrization of Katz et al. (2003). The second condition arises from the top of the




Figure 5. Gravity anomalies expected at the Earth's surface when the lithospheric thickness is 100 km. Plots are as in Figure 3, except the attenuation of gravity anomalies through the mechanical boundary layer, and an elastic plate with $T_e = 30$ km has been taken into account.

convecting system not being at the Earth's surface. We envisage that there is a rigid mechanical boundary layer (MBL) which separates the Earth's surface from the top of the convecting system. The temperature near the base of the MBL is where T/T_s (where T_s is the melting temperature) is highest and therefore the viscosity is lowest (e.g., Frost & Ashby, 1982). For this reason we used both free-slip and rigid boundary conditions on the top surface of the convecting box. A stress-free boundary is probably the better approximation to the behavior of the real Earth. Heat transfer through the MBL is purely by conduction, and we assume the thermal profile is linear through this region and equal to 0°C at the Earth's surface. At the top of the convecting region we assume that both the horizontally averaged temperature and heat flux are continuous with that in the MBL. Finally, we determine the thickness of the MBL by prescribing a lithospheric thickness, which we define as the intersection of the linear conductive profile of the MBL with an isentropic




Figure 6. Topography expected at the surface, after flexural filtering through an elastic plate with $T_e = 30$ km.

profile at the interior mantle potential temperature. In what follows the lithosphere thickness is fixed at 100 km, except when considering melt generation where we have varied this parameter, as melt generation is particularly sensitive to it. This choice of interior potential temperature and lithospheric thickness fixes the heat flux for all simulations to be 50 mW m^{-2} , which is similar to that through old sea floor (Hasterok, 2013). The resulting associated parameters, which include the upper mantle viscosity, are given in Table 2. The horizontally averaged potential temperature profiles after scaling are shown in Figure 2.

With Rayleigh numbers in the range 10^5 to 10^6 , Table 2 shows mantle viscosity values vary from 1.4×10^{20} to $5.3 \times 10^{21} \text{ Pa s}$, which is around the range expected for the upper mantle ($\sim 10^{21} \text{ Pa s}$). The higher the Rayleigh number, the lower the inferred viscosity, the smaller the potential temperature difference across the layer, and the thicker the MBL. Section A4 discusses further the behavior of the parameters with Rayleigh number.




Figure 7. Cross plots of gravity against topography at the top of the convecting box for the four $Ra = 10^6$ simulations. In each case a geometric regression line has been calculated, and marked on each plot are the slope Z of that line and r^2 , the square of the correlation coefficient. If the boxes are overlain by water, the values of Z are reduced by a factor of $(\rho_0 - \rho_w)/\rho_0 \approx 0.7$.

3. Gravity and Topography

Examples of the gravity and topography at the top of the convecting box when there is no lithosphere present are shown in Figures 3 and 4, calculated from the expressions in Parsons and Daly (1983) (section A1). The gravity field takes account of the contribution from the thermal expansion of the fluid and that from the topography (assuming deformable top and bottom boundaries). Both shadowgraphs and the plots of gravity and topography are filtered versions of the temperature field in the box, but they involve distinctly different kinds of filters. Since the shadowgraph represents the Laplacian of the vertically averaged temperature, it is equally sensitive to temperature variations at all depths and acts to emphasize short-wavelength features. By contrast, gravity and topography act to attenuate short-wavelength features in the temperature field, particularly those at depth (Parsons & Daly, 1983).

Before comparing the results from the numerical models with the observations a further filter needs to be applied, to account for the effect of the overlying mechanical boundary layer. This layer has a number of effects. First, gravity anomalies are attenuated by the thickness of the layer. Second, the elastic properties of the overlying plate acts to filter out the topographic expression of short-wavelength features, with the magnitude of this effect dependent on the assumed effective elastic thickness (T_e). Figures 5 and 6 show the expected gravity and topography at the surface, after applying a filter for the mechanical boundary layer, assuming an elastic thickness $T_e = 30$ km. The effect of all this filtering is to remove much of the short-wavelength information that is visible in the shadowgraph images. In particular, the filters emphasize




Figure 8. (a) Power spectral density (PSD) of gravity anomalies at the top of the convecting region as a function of wave number $k = 2\pi/\lambda$, for the four convection simulations at $Ra = 10^6$, along with that estimated for the Earth from the gravity model DIR-R5. Log scales are used for both axes. The approximate spherical harmonic degree estimated using Jeans relation is plotted along the top axis. Two thin line segments show slopes which yield a power law decay proportional to k^{-2} (Kaula's rule) and k^{-3} . (b) Power spectral density of dynamic topography at the top of the convecting box as a function of wavenumber, along with that estimated for the Earth by Hoggard et al. (2016). (c) Admittance (the ratio of gravity to topography) at the top of the convecting region as a function of wave number, assuming air loading. (d) Coherence (square of the correlation between gravity and topography) at the top of the convecting region as a function of wave number.

the “hubs” of the convective pattern and suppress the expression of the “spokes.” As will be seen in the next section, the effect of the filter on melt generation is even more important. As the thickness of the MBL increases it suppresses melt generation first in the spokes and then in the hubs.

The images in Figures 5 and 6 demonstrate three key points: First, the predicted amplitude of gravity anomalies is comparable with the observed anomalies (compare the scale of Figure 5 with that of Figure 13c). Second, the spatial patterns of gravity anomalies and topography are sensitive to the assumed boundary conditions. Third, there is a good correlation between the topography and the gravity in plots of the values of gravity and topography at each spatial location (Figure 7), which closely resemble similar plots that have been made using observed values of gravity and residual topography (e.g., see Figures 6 and 7 of Crosby & McKenzie, 2009). The slope of these plots represents a characteristic average value of the admittance (ratio of gravity to topography) associated with the convection. A characteristic air-loaded admittance between 43 and 53 mGal km^{−1} is inferred from Figure 7, corresponding to values of 30–37 mGal km^{−1} when overlain by water. These values are similar to those previously reported (Parsons & Daly, 1983).

An alternative way to assess the predictions of gravity and topography is to work in the frequency domain rather than the spatial domain. Figures 8a and 8b show the power spectral density of gravity and topography at the top of the convecting box (section A2), along with estimates for the Earth. The figures show that the

convection experiments have most power over a range of wavelengths from around 500 to 2,000 km, where the estimates of gravity anomalies are comparable in magnitude with those of the Earth. This behavior reflects the fact that the convection organizes into cells where the distance from one upwelling to the next is around twice the layer depth, corresponding to a wavelength around 1,200 km. Away from this broad peak the power from the convection experiments decays. At the short wavelengths shown in the plot, this decay in the power spectral density scales roughly as k^{-3} . The different boundary conditions in the convection experiments lead to subtle differences in the gravity spectra in Figure 8. For example, there is greater power at long wavelengths for those simulations with free-slip bottom boundary conditions than rigid. There is also greater power at short wavelengths for the simulations with rigid top boundary conditions than free-slip top boundary conditions.

The power spectral density of gravity for the Earth is notably different from that of the convection experiments. Overall, the Earth's power spectral density decays broadly with wavenumber as k^{-2} , a power law which is referred to as Kaula's rule. In detail, the slope of decay flattens slightly in the 2,000 to 500 km wavelength band, before becoming steeper at wavelengths shorter than 500 km, but not decaying as steeply as in the convection experiments. The difference between the convection experiments and the Earth for wavelengths shorter than 500 km is to be expected: On Earth surface loading causes short-wavelength topography that is supported by elastic stresses in the plate. The short-wavelength power in the Earth's gravity field arises from surface loading, not mantle convection. At long wavelengths ($>2,000$ km) mantle convection should play an important role in determining the gravity field on Earth, but the convection experiments here have notably less power than the Earth. Thus, the upper mantle convection modeled here does not account for the magnitude of the long-wavelength portion of the Earth's gravity field.

Figure 8b shows the corresponding spectra for predictions of dynamic topography. The shape of the topography spectra of the convection runs is different from that for the gravity. The topography spectra are broadly flat up to around a wavelength of 1,000 km and then decay steeply at shorter wavelengths. Thus, dynamic topography has relatively more long-wavelength power than does gravity, and this can be seen in the space domain plots of Figures 3 and 4—the topography plots look smoother than the gravity plots (as noted by Craig & McKenzie, 1987).

It is more difficult to compare predictions of dynamic topography with observations. The reason for this is that there are significant long-wavelength features in the Earth's topography that are not associated with mantle convection, for example, the difference in elevation between the oceans and the continents, and other topography associated with variations in crustal thickness. A recent attempt has been made to estimate the power spectrum of dynamic topography by Hoggard et al. (2016), by fitting spherical harmonics to point observations of residual depth in the oceans, and a fixed scaling of long-wavelength gravity anomalies in the continents. Their estimate of power indicates a Kaula-rule-like decay in the power spectral density as k^{-2} and is plotted in Figure 8b. At wavelengths longer than 2,000 km the convection experiments are not able to explain the Hoggard et al. (2016) estimate of dynamic topography. At shorter wavelengths spectra become more comparable, although only a limited comparison can be made because the Hoggard et al. (2016) estimate is limited to spherical harmonic degree 30 (a wavelength of 1,300 km). The Hoggard et al. (2016) estimates are based on air loading in the continents and water loading in the oceans. The power spectra of the convection experiments plotted in Figure 8b assume air loading. The effect of water loading would be to increase the amplitude of the power spectral density of the topography by a factor of $(1 - \rho_w/\rho_0)^2 \approx 2$, which would represent only a small shift on the log-scale plot of Figure 8b.

In addition to comparing the observed and calculated individual spectra of gravity and topography, it is also of interest to look at their relationship to one another (Parsons & Daly, 1983). This relationship is typically characterized in terms of the admittance and coherence of the two signals, with the topography taken as input and the gravity as output (section A3). Figure 8c shows the admittance in the spectral domain, calculated from the numerical experiments. The value of the admittance increases with increasing wave number (decreasing wavelength), approximately as $\log(k)$ for the range of wave numbers shown. The behavior of admittance with wave number is similar for the different boundary conditions over the range of interest, with significant departures only noticeable at long wavelengths ($>2,000$ km). Indeed, at long wavelengths the admittance for the free-rigid case becomes negative around a wavelength of 3,700 km (as noted by Parsons & Daly, 1983, see their Figure 5). Figure 8c is consistent with the behavior in the cross plots of Figure 7. The gravity anomalies have most power at wavelengths around 1,000 km, corresponding to an air-loaded

admittance of around 45 mGal km^{-1} in Figure 8c, which is broadly the slope obtained from the cross-plots. The corresponding coherence in Figure 8d is close to 1, reflecting the good correlation between the two observables that can be seen in the space domain plots. Only at wavelengths longer than 2,000 km is a weak coherence between the signals seen, and then only significantly for the free-rigid case. This behavior mirrors the admittance at long wavelengths for the free-rigid case, where gravity and topography correlate positively except for wavelengths longer than 4,000 km when they correlate negatively.

The frequency domain plots in Figure 8 illustrate the spectral properties of the gravity and topography at the top of the convecting box, as shown in the space domain in Figures 3 and 4. When considering signals at the Earth's surface, the spectral properties will be further modified by the filtering effect of the MBL on top. How significant this effect is depends on the effective elastic thickness and the thickness of the MBL. For an elastic thickness $T_e = 30 \text{ km}$ as used in Figures 5 and 6, the wavelength at which the Fourier coefficients of topography are reduced by a factor of 2 is $\lambda_{\text{flex}}^{1/2} = 330 \text{ km}$. Wavelengths shorter than this will be significantly attenuated by the flexural filtering; wavelengths longer than this will not. The topography spectra in Figure 8 would only be significantly different at wavelengths shorter than $\lambda_{\text{flex}}^{1/2}$ were a flexural filter to be applied. The effect of flexural filtering on the gravity is more complicated and acts to produce a modest increase in the gravity signal in a wavelength band around that associated with the MBL thickness and that associated with flexure (see section A1 and Figure A1 for further discussion).

4. Melt Generation

The generation of melt, its separation from its source regions, and the time τ required for it to move from its source to the surface have all been extensively studied, both theoretically using two-phase flow equations and observationally using a variety of geochemical approaches. The two-phase flow equations show that basaltic melt separates from its source regions when the melt fraction by volume ϕ_0 exceeds $\sim 0.5\%$ (McKenzie, 1985b). Studies of the composition of abyssal peridotites (Johnson et al., 1990; Warren, 2016) show that the incompatible elements that were present in the source before melting occurred have been removed by the melt. Estimates of the melt fraction present during the melting are between 0.2% and 0.7% (Liang & Peng, 2010; Slater et al., 2001). Estimates of ϕ_0 and τ can also be obtained from measurements of U series disequilibria (Kokfelt et al., 2003; Koornneef et al., 2012; McKenzie, 1985a; Stracke et al., 2006; Turner et al., 2016). Most estimates give $\phi_0 \leq 0.5\%$ and $\tau \leq 1 \text{ ka}$. Perhaps the strongest constraint on the values of both ϕ_0 and τ comes from modeling the generation of melt by deglaciation. When most of the ice covering Iceland melted at the end of the last glaciation the melt production rate suddenly increased (Eason et al., 2015; MacLennan et al., 2002). The thickness of the ice that melted was about 2 km, increasing the melt fraction present in the source region by only about 0.2% (Eksinhol et al., 2019; Jull & McKenzie, 1996). This increase was sufficient to generate large shield volcanoes within about 1 ka of the removal of the ice. These models and observations all show that melt generated by decompression melting in the upper mantle rapidly moves to the surface and that no appreciable volume remains in the source region. We therefore calculated the rate of melt production by simply vertically integrating the melting rate over the thickness of the layer in which melt was being produced.

Figure 9 shows the calculated rate of melt production for the four $Ra = 10^6$ simulations, with a lithospheric thickness of 80 km, and 100 ppm water. The hydrous melting parametrization of Katz et al. (2003) was used (section A5). Melt production in Figure 9 only occurs where the mantle is upwelling, where the gravity anomalies and topography are positive (Figures 5 and 6). However, the spatial extent of the regions of high melt rate are smaller than the regions of positive gravity anomaly. Most of the melting takes place in the regions directly above the narrow ($\sim 60 \text{ km}$ diameter) upwelling plume conduits, where the decompression rate is greatest. However, melting also takes place, albeit to a lesser degree, in a broader region around the conduits and above some of the rising sheets connecting neighboring plumes (i.e., on the spokes as well as the hubs of the spoke pattern). Melt generation along some of the spokes is particularly clear for the free-free simulation, with linear bands of relatively low melt production connecting concentrated centers of relatively high melt production.

The rate of melt production is particularly sensitive to the thickness of the lithosphere and the water content of the mantle. Figure 10 illustrates the effect of varying these two parameters for the free-rigid $Ra = 10^6$ simulation. At low water contents, and beneath thick lithosphere, melting is restricted to the hubs in the spoke pattern, if indeed melting happens at all. However, high water contents and thin lithosphere result in




Figure 9. Vertically integrated rate of melt production beneath an 80 km thick lithosphere with 100 ppm water. Note that the color scale for melt rate is logarithmic. Numbers in white give the total rate of melt production in $\text{km}^3 \text{ ka}^{-1}$ for each contiguous zone of melting. Regions of total melt production less than $1 \text{ km}^3 \text{ ka}^{-1}$ are not labeled.

melting along the spokes. As Figure 10 illustrates, for lithosphere as thick as 180 km, melting is suppressed unless the water content is sufficiently high ($\sim 1,000$ ppm). For lithosphere as thin as 70 km, melting along both the hubs and the spokes can be seen even for modest (~ 100 ppm) water contents. The water content of the convecting upper mantle is probably between 100 and 200 ppm (Michael, 1995; Saal et al., 2002). In contrast, the water content is likely to be considerably greater where the base of the lithosphere has been enriched by metasomatism. It is not straightforward to estimate the water content of such regions using that in the nodules brought up by magmas such as kimberlites from depths of 100–200 km, because they are often infiltrated by the host magma. Protons are especially mobile. More reliable estimates can be obtained from the Ce concentration, because Ce and H have similar bulk partition coefficients between magma and peridotite (Aubaud et al., 2004), and the Ce concentration in nodules is less affected by infiltration than is




Figure 10. Vertically integrated rate of melt production for the free-rigid simulation with $Ra = 10^6$, showing the effect of varying the water content (as labeled horizontally in ppm), and different lithospheric thicknesses (as labeled vertically in km).

that of H (Erlank et al., 1987 p 283). The Ce concentration in the commonest class of nodules is ~ 10 ppm (Erlank et al., 1987), or about 10 times that of the convecting upper mantle. Therefore, the metasomatically enriched region at the base of thick old lithosphere probably has a water concentration of $\sim 1,000$ ppm. Where the lithosphere is thin, Figure 10 shows that such high water concentrations will lead to widespread melting along spokes, and at the hotter hubs melting at depths as great as 180 km.

5. Terrestrial Observations

The numerical experiments described above show that the planform of mantle convection will be most obviously expressed in the surface observables when both the elastic thickness and lithospheric thickness are small. Figure 11 shows that the lithospheric thickness exceeds 120 km over large regions of western and southern Africa. In these regions the volcanism consists of small-volume alkalic eruptions, such as kimberlites, that contain high concentrations of carbonates and hydrous minerals. Where kimberlites are diamond-bearing, Figure 11 shows that the lithospheric thickness generally exceeds 150 km. The limited




Figure 11. Lithospheric thickness calculated from surface wave tomography (Priestley et al., 2018). The thick square box indicates a region that is of the same horizontal extent as the convection simulations. The elastic thickness of the region within the dotted lines is likely to be less than 4 km (Figure 12).

spatial resolution of surface wave tomography, of ~ 250 km, probably accounts for the few diamond-bearing locations in Figure 11 that appear to have thinner lithosphere.

The lithospheric thickness within much of the rectangle marked by the thick continuous black line in Figure 11 is less than 80 km, and the horizontal extent of this region is similar to that of the numerical experiments. In the eastern and northern parts of the area there are sufficient surface gravity measurements to allow T_e to be estimated from the transfer function between the free air gravity anomalies and the topography (McKenzie & Fairhead, 1997), giving values of 3–4 km (see supporting information). The coherence method and Bouguer anomalies provides only an upper bound, not an estimate, of the value of T_e (McKenzie, 2016). Within the box marked by the dotted lines in Figure 11 there are few surface gravity measurements. Instead the satellite gravity field DIR-R5 can be used to show that the admittance is about 50 mGal km^{-1} between wavelengths of 200 and 1,000 km, and that the elastic thickness is probably less than 4 km (Figures 12a and 12b). Therefore, in this region and wavelength band both the gravity and topography are controlled by convection. The same is not the case in southern Africa, where the elastic thickness is about 30 km (McKenzie et al., 2015).

Figure 13 shows maps of gravity, topography, and subaerial volcanism across Africa and Arabia. The volcanism beneath the Red Sea and in the Afar results from upwelling of the mantle between separating plates.

However, in Ethiopia, Kenya, and Kivu the upwelling from the limited extension is insufficient to account for the extensive volcanism. There is no obvious orientation of the topography and gravity anomalies, probably because Africa is almost stationary with respect to the hot spot frame. The correspondence between the features in Figure 13 and the maps from the numerical experiments is striking. Volcanism is almost entirely restricted to regions where the lithospheric thickness is less than 70 km. The only clear exception is Kivu, where the spatial resolution of the surface wave tomography is probably insufficient to resolve the thickness of the thin lithosphere beneath the Western Rift. The linear volcanic feature extending from Kenya to the Kars Plateau resembles similar linear features in Figure 9, with localized regions of concentrated upwelling being associated with positive gravity and elevation, and with enhanced volcanism. Like the numerical experiments, the volcanism is more localized than are the associated positive gravity and topographic anomalies. What is less clear is which of the four combinations of boundary conditions best fits the observations. Figures 5 and 6 show that the observed horizontal scales of the convective features are probably larger than those of the rigid-rigid, and smaller than those of the free-free, experiments. The scales of the anomalies in the other two experiments are similar, both to each other and to the observed scales. The rigid-free experiment has broadly circular patterns of positive gravity anomalies surrounded by linear negative anomalies, whereas the free-rigid case has the opposite. The viscosity of the lower mantle is greater, and that of the asthenosphere immediately below the lithosphere less, than that of the upper mantle. These viscosity variations suggest that the free-rigid experiment should match the observed patterns better than the rigid-free case. However, the patterns in Figure 13 are not obviously more like the free-rigid features than the rigid-free ones. Furthermore, the variation of viscosity with temperature, which has been ignored, may have a strong influence on the geometry and in particular whether the planform is dominated by rising or sinking fluid in the hubs. At low Rayleigh numbers the answer to this question is controlled by the sign of $d\eta/dT$, with the flow in the plumes being in the direction of increasing viscosity (Segel & Stuart, 1962). If the same is true at large Rayleigh numbers the hubs in both these experiments will consist of hot rising material.

The Cameroon Line forms a curve, similar to features in Figure 9. As the lithosphere thickness increases to the SW, where the Line lies beneath Atlantic lithosphere, the volcanism decreases. The association of positive gravity anomalies, elevated topography, and volcanism is clearly expressed even in relatively small features like Air and Darfur. Many volcanic centers, such as Air and Hoggar, and Haruj and Tibesti, are linked to each other by lines of positive gravity and topography, where there is limited volcanism. An especially obvious feature extends from South Arabia to Anatolia, where the volcanism is beneath Western Arabia, not the Red Sea. Such lines are most clearly visible in Figures 13c and 13e where the lithosphere is thin beneath NE




Figure 12. Plots of the admittance and coherence of gravity and topography versus $1/\lambda$ where λ is the wavelength and $k = 2\pi/\lambda$ the wavenumber, in three regions. (a) and (b) use free air gravity anomalies and topography from inside the box in Figure 11 marked by the dotted line. The gravity field was calculated from DIR-R5 (Bruinsma et al., 2014) by setting the coefficients from $l = 2$ to 7 to 0 and applying a taper $f = (l - 7)/5$ to those from $l = 8$ to 11 (wavelengths 3,333 to 5,000 km). A low-pass filter falling to 1/2 at 250 km was applied to remove the short-wavelength anomalies associated with elastic flexure. The topography is taken from ETOPO5. The admittance was calculated using the topography as input, gravity as output. The two dotted lines show the flexural admittance for two values of the elastic thickness T_e . (c)–(f) show corresponding plots for the Pacific (see supporting information for maps) and Indian Oceans (see Figure 14). The gravity anomalies were calculated from the DIR-R5 coefficients with those of degree 2 set to 0. The admittances in (c) and (e) were calculated from the ratio of the spectral coefficients. The dotted lines show the flexural admittance for $T_e = 20$ km.




Figure 13. Maps of the volcanism (a), lithospheric thickness (b), gravity (c and d), and topography (e and f) within the region marked by the heavy black line in Figure 11. The gravity and topography are shown without (c and e) and with (d and f) the regions covered by volcanics, most of which are Miocene or younger, taken from Thorpe and Smith (1974), Ball et al. (2019), and earthwise.bgs.ac.uk.

Africa, Arabia and Anatolia. All these features are similar to those of the numerical experiments, and all are consistent with a spoke pattern of convection existing beneath the region. In particular, and as expected from Figure 9, the extent of the volcanism is controlled by variations in lithospheric thickness and is limited in the south and east by the thick lithosphere of the Congo Craton.

The Cameroon Line and the line of active volcanism that extends from Afar to eastern Anatolia have long puzzled geophysicists, because their geometry is difficult to reconcile with a planform of mantle convection consisting of plumes. Sleep (1997, 2008) and Ebinger and Sleep (1998) argued that these linear volcanic features were produced by lateral flow from plumes in channels beneath the lithosphere. One problem with this proposal is that the composition of the volcanics along the Cameroon Line shows so little variation (Fitton, 1987; Lee et al., 1994).

A different model was proposed by Milelli et al. (2012) which emphasized the location of the volcanism, which has remained in the same region of Africa as the continent has moved. They argued that this behavior required the volcanism to result from thermal instabilities in the lower part of the lithosphere, rather than being the surface expression of convective upwellings in the upper mantle below the plates. The numerical experiments described above show that the expected planform of upper mantle convection is that of hubs joined by spokes, both of which can generate melt if the lithosphere is sufficiently thin. The observed linearity of the Cameroon Line and other features in NE Africa therefore requires no special explanation. The experiments also show that Milelli et al. (2012)'s observations may also have a simple explanation, since the melting rate, and not the planform, is controlled by the lithospheric thickness, and the volcanism of the Cameroon Line lies along the northern edge of the thick lithosphere of the Congo Craton. In contrast the line of volcanism from the Afar to Kars lies within a region of relatively uniform lithospheric thickness. It is therefore unlikely that such linear features all form from edge convection like that discussed by King and Anderson (1998).

In general, it is not possible to compare the melt generation rates in Figure 9 with those observed because they are so rarely estimated by the geologists who map the volcanics. An exception is Mount Cameroon, which is the most active volcano in Africa. Its eruption rate was estimated by Suh et al. (2003) to be about $700 \text{ km}^3/\text{Ma}$. When the lithospheric thickness is 80 km the larger hubs in the free-rigid experiment in Figure 9 produce about $6 \times 10^4 \text{ km}^3/\text{Ma}$ and the smaller ones $1 \times 10^3 \text{ km}^3/\text{Ma}$. The rates of melt generation in the numerical experiments can therefore easily account for the observed rates. But they are quite inadequate to account for the production rates that occur during major flood volcanism, which commonly exceed $1 \times 10^6 \text{ km}^3/\text{Ma}$. Like the long-wavelength gravity anomalies discussed below, simple isoviscous upper mantle convective models cannot account for such events.

The box in Figure 11 is too small to be used to study the long-wavelength components (wavelengths $> 1,000$ km) of the Earth's dynamic topography and gravity. These are best studied in the Indian and Pacific Oceans. The elastic thickness of old oceanic lithosphere is about 20 km (e.g., McKenzie et al., 2014). Therefore, wavelengths greater than about 800 km are little affected by the thickness and elastic properties of the lithosphere (Figures 12c and 12e). In many continental regions the surface topography is dominated by isostatically compensated variations in crustal thickness. Such structures are less common in oceanic regions, where the bathymetry is principally controlled by the age of the lithosphere. The effect of plate cooling can be removed by using a depth-age model, and regions of thick crust removed by hand (e.g., Crosby et al., 2006; Crosby & McKenzie, 2009; Hoggard et al., 2017). The resulting residual topography is then largely supported by convection. Figure 12 shows a comparison between free air gravity, calculated from DIR-R5 with the coefficients of degree 2 set to 0, and Crosby's values of residual depth in the Indian and Pacific Oceans. Those obtained from Hoggard's values of residual depth are similar and are shown in the supporting information. Figure 14 shows maps of such anomalies in the Indian Ocean: Those for the Pacific are illustrated in the supporting information. The admittance between wavelengths of 1,000 and 2,000 km is about 30 mGal km^{-1} , in agreement with the values from the numerical experiments in Figure 8. However, at wavelengths greater than about 2,000 km the gravity and residual depth anomalies in both oceans cease to be coherent. This incoherency is particularly striking in the Indian Ocean, where the large negative gravity anomaly covering the NE part of the Ocean (Figure 14a) has no expression in the residual depth (Figure 14b). At wavelengths greater than 2,000 km the observed power spectrum also differs from that calculated from the box models (Figure 8a). The power in the observed gravity field continues to increase at wavelengths longer than 2,000 km, unlike that from the numerical experiments. This behavior shows that simple isoviscous convective




Figure 14. Gravity and residual depth in the Indian Ocean. (a) Gravity from DIR-R5, with coefficients $l = 2$ set to 0 and a filter applied, falling to $1/2$ at 250 km, to remove the short-wavelength components. (b) Residual depths, averaged over $2^\circ \times 2^\circ$ boxes (Crosby et al., 2006). The dots show the locations of the resulting averages. Airy projection with center 30°N , 80°E . $\beta = 30^\circ$.

models cannot account for the longest wavelength part of the Earth's gravity field, and suggest that it is not maintained by upper mantle convection. Though the size of the boxes used for the numerical experiments is too small to determine the power at such wavelengths accurately, there is no suggestion in the planforms that the presence of lateral boundaries governs the scale of the convection.

6. Conclusions

The numerical experiments described above show that the observed gravity and topographic anomalies are reproduced by the simplest isoviscous fluid dynamical model of thermal convection. The wavelength at which convective support dominates elastic support is controlled by the elastic thickness T_e and varies from about 200 km where $T_e \leq 4$ km in NE Africa to ~ 500 km in southern Africa, where $T_e \sim 30$ km. Melt generation occurs where mantle material moves upward and is therefore controlled by the lithospheric thickness

and not by the value of T_e . The correspondence between the volcanism and the gravitational and topographic anomalies in NE Africa is striking and shows that they all result from the convective circulation.

There is no similar correspondence between the results from the numerical experiments and the gravity and residual depth anomalies at wavelengths greater than 2,000 km. Furthermore, the absence of correlation between residual depth and gravity anomalies with wavelengths greater than 2,000 km in the Pacific and Indian Oceans is unlike the behavior observed at shorter wavelengths in these oceans. The simple numerical models discussed here cannot account for the long-wavelength gravity anomalies with spherical harmonic degrees $l \leq 20$.

The close correspondence between the calculated and observed topography, gravity, and volcanism suggests that it should be possible to use the surface observables where the lithosphere is thin and T_e is small, together with the isoviscous convective equations, to map the convective circulation in the upper mantle.

Appendix A: Numerical Methods

A1. Gravity, Topography, and Flexure

To calculate surface gravity and topography, the second-order finite element temperature fields were first linearly interpolated onto a regular grid of points with 64 mesh points in the vertical and 512 in the horizontal. The method of Parsons and Daly (1983) and Craig and McKenzie (1987) was then used to calculate both topography and gravity, by Fourier transforming the grids of data, multiplying by an appropriate filter (given in Appendix A of Parsons & Daly, 1983, for the different boundary conditions), and transforming back. For gravity calculations both the top boundary and the bottom boundary were assumed to be deformable. The deformation of the top and bottom boundaries was calculated using the approximation $(\rho_0 - \rho_w)gh = -\sigma_{zz}$, where h is the deformation of the interface, σ_{zz} is the normal stress at the relevant boundary in the fixed geometry of the convection simulations, and ρ_w is the density of the fluid on the other side of the boundary.

The effect of an elastic layer above the convecting fluid is to low-pass filter the dynamic topography (Figure A1b). To produce the flexurally filtered topography seen in Figure 6, the Fourier coefficients of the topography in Figure 4 were multiplied by the Fourier-domain flexural filter,

$$F(k) = \frac{1}{1 + (\alpha_{\text{flex}} k)^4}, \quad (\text{A1})$$

where k is the wave number, and the flexural parameter α_{flex} is related to the elastic thickness T_e by

$$\alpha_{\text{flex}} = \left(\frac{ET_e^3}{12(1 - \nu^2)(\rho_0 - \rho_w)g} \right)^{1/4}. \quad (\text{A2})$$

The Fourier coefficients are reduced by a factor of 2 at a characteristic wavelength $\lambda_{\text{flex}} = 2\pi\alpha_{\text{flex}}$. Values of the Young's modulus $E = 10^{11}$ Pa and Poisson's ratio $\nu = 0.25$ are assumed, such that $\lambda_{\text{flex}}^{1/2} = 330$ km for an elastic thickness $T_e = 30$ km; and $\lambda_{\text{flex}}^{1/2} = 85$ km for an elastic thickness $T_e = 5$ km. All plots assume air loading and thus set $\rho_w = 0$. For water loading $\rho_w = 1,000 \text{ kg m}^{-3}$.

The effect of an elastic layer on gravity anomalies is subtle, because there is a contribution to gravity anomalies both from the topography and from the density variations at depth, which will be attenuated by a factor e^{-kt_m} if there is a MBL of thickness t_m on top. The gravity anomalies in Figure 5 are shown after this filtering process, which is calculated assuming the MBL is laterally uniform. The corresponding spectra are shown in Figure A1a. If Δg_0 and h_0 represent the gravity and topography estimated at the top of the convecting box, the corresponding gravity Δg_1 at the surface can be calculated in the Fourier-domain from

$$\Delta g_1(k) = (\Delta g_0(k) - 2\pi G \rho_0 h_0(k)) e^{-kt_m} + 2\pi G \rho_0 h_0(k) F(k), \quad (\text{A3})$$

assuming air loading. This equation splits the gravity anomaly into two components: The term on the far right represents the gravity due to the surface topography, which is attenuated according to the factor $F(k)$ in (A1). The other term represents the density variations at depth, which are attenuated as e^{-kt_m} due to the finite thickness of the MBL. At long wavelengths gravity anomalies are unchanged by this additional filtering. At wavelengths significantly shorter than both the flexural wavelength λ_{flex} and the wavelength $\lambda_m = 2\pi t_m$ associated with the MBL, gravity anomalies are strongly attenuated, as both the e^{-kt_m} and $F(k)$ filters




Figure A1. Spectral characteristics as in Figure 8, but at the surface after filtering through the MBL and an elastic layer with thickness $T_e = 30$ km. Notice the significant decrease in the power of gravity anomalies at short wavelengths and a moderate increase in the power at intermediate wavelengths.

tend to 0. At intermediate wavelengths, particularly in the wavelength band around λ_m and $\lambda_{\text{flex}}^{1/2}$, gravity anomalies actually slightly increase in magnitude due to this additional filtering. This behavior occurs because a MBL acts to separate mass excesses at the surface (associated with positive gravity anomalies) from mass deficits at depth (associated with negative gravity anomalies). The thicker the MBL, the greater the attenuation of the negative anomalies, and the larger the net positive anomaly. Correspondingly, there is a modest increase in the admittance due to the addition of a MBL (Figure A1c; see section 2.3 of Crosby & McKenzie, 2009, for further discussion). It should be noted that having a laterally uniform thermal structure in the MBL is a poor approximation, as it implies a discontinuity in heat flux between the MBL and the top of the convecting box (where the heat flux varies laterally). However, we have made such an approximation here because the convection simulations fix the temperature at the top of the convecting layer, rather than at the Earth's surface. A better approach would model the temperature structure of the MBL during the convection simulations. However, such a modification is unlikely to make more than minor changes to the results of the calculations, because the thickness of the MBL is small compared to that of the convecting layer (Table 2) and the temperature on the upper boundary of the MBL is fixed. If the temperature structure of the MBL is included in the calculations it is then no longer accurate to obtain the topography and gravity for variable thicknesses of the MBL simply by filtering in the spectral domain.

A2. Power Spectral Density

Power spectral density (PSD) estimates were calculated using the method described by Rexer and Hirt (2015). The initial data (either gravity or topography) is a regularly spaced grid of points representing a region of dimensional extent L_x by L_y . Let the number of grid points in the x direction be N , and the number

in the y direction be M . The initial data are given as a matrix of values d_{rs} where $r = 0, 1, \dots, N-1$, and $s = 0, 1, \dots, M-1$. The convection simulations have reflecting boundary conditions at the sides, so the natural Fourier representation to use is a discrete cosine transform of the first type (DCT-I), which is equivalent to a discrete Fourier transform of a $2N-2$ by $2M-2$ extended grid of data exploiting the even symmetry. The discrete Fourier coefficients f_{pq} are defined by equation (8) of Rexer and Hirt (2015),

$$f_{pq} = \frac{1}{(2N-2)(2M-2)} \sum_{r=0}^{2N-2} \sum_{s=0}^{2M-2} d_{rs} \exp\left(-\pi i \left(\frac{sp}{N-1} + \frac{rq}{M-1}\right)\right). \quad (\text{A4})$$

The even symmetry extends the data such that for $N \leq r \leq 2N-2$ the value is taken from the original grid at $r' = 2N-2-r$ and for $M \leq s \leq 2M-2$ the value is taken from original grid at $s' = 2M-2-s$. In dimensional variables, the corresponding wavenumbers of the transform are $k_p^x = \pi p/L_x$ and $k_q^y = \pi q/L_y$. Owing to the reflection boundary conditions, the discrete Fourier coefficients f_{pq} are real and even in both directions. A 2-D grid of power spectral density can then be computed from equation (10) of Rexer and Hirt (2015),

$$\phi_{pq} = 4L_x L_y |f_{pq}|^2. \quad (\text{A5})$$

Finally, the 2-D-PSD were then azimuthally averaged in wavenumber space with a bin width of $1.3 \times 10^{-3} \text{ km}^{-1}$, to produce the 1-D-PSD profiles that are shown in Figures 8a and 8b (with units of either $\text{mGal}^2 \text{ km}^2$ or $\text{km}^2 \text{ km}^2$).

Data for the Earth are typically given in terms of spherical harmonic coefficients, which need to be manipulated before they can be compared directly with the 1-D-PSD profiles calculated for the Cartesian geometry of the convection simulations. This process is also described by Rexer and Hirt (2015). The spherical harmonic degree l can be related to the Cartesian wavenumber k by the Jeans relation approximation

$$k = \frac{l + \frac{1}{2}}{R} \quad (\text{A6})$$

where R is the radius of the Earth. An estimate of the 1-D Cartesian PSD can be obtained from

$$\phi_{\text{PSD}}(k) = 4\pi R^2 \frac{P_l}{2l+1} \quad (\text{A7})$$

where P_l is the power at spherical harmonic degree l (see equation (13) of Rexer & Hirt, 2015).

A3. Admittance and Coherence

The admittance in Figure 8c was computed as

$$Z(k) = \frac{\langle \Delta g, h \rangle}{\langle h, h \rangle} \quad (\text{A8})$$

where $\langle \cdot, \cdot \rangle$ represents the cross-power of the signals as a function of wavenumber k , calculated in the same way as the power spectra by multiplying the Fourier coefficients and then azimuthally averaging. Since the Fourier coefficients are real, the admittance is also real. The coherence in Figure 8d was calculated similarly as

$$\gamma^2(k) = \frac{\langle \Delta g, h \rangle^2}{\langle \Delta g, \Delta g \rangle \langle h, h \rangle}. \quad (\text{A9})$$

A4. Rayleigh Number Scalings

Boundary layer theory suggests that there should be systematic power law scalings with Rayleigh number for properties of the convecting system, such as the thickness of boundary layers, and the heat flux. Table A1 shows such approximate scaling laws that have been obtained from the 12 numerical runs presented here. These scaling laws should be used with caution: They were determined from the properties of the system at a single snapshot in time, for a limited range of Rayleigh numbers. However, they illustrate the broad trends. As expected, Nusselt number increases with Rayleigh number: The behavior for the free-free system as $\text{Nu} \propto \text{Ra}^{0.3}$ is close to the one-third power law expected from boundary layer theory (McKenzie et al., 1974). Since the dimensional scaling used in the main text essentially fixes the heat flux, the potential temperature difference across the layer given in Table 2 scales as the inverse as the Nusselt number scaling, that

Table A1

Approximate Scalings with Rayleigh Number for Nu, the Nusselt Number (Ratio of Convective to Conductive Heat Transfer); $h'_{\text{RMS}} = h_{\text{RMS}} / (\rho_0 \alpha \Delta T_p d / (\rho_0 - \rho_w))$, Dimensionless Root-Mean-Square Dynamic Topography; and $\Delta g'_{\text{RMS}} = \Delta g_{\text{RMS}} / (2\pi G \rho_0 \alpha \Delta T_p d)$, Dimensionless Root-Mean-Square Gravity Anomaly

BC	Nu	h'_{RMS}	$\Delta g'_{\text{RMS}}$
Rigid-rigid	$0.189 \text{ Ra}^{0.281}$	$1.682 \text{ Ra}^{-0.342}$	$0.405 \text{ Ra}^{-0.308}$
Free-rigid	$0.228 \text{ Ra}^{0.283}$	$1.098 \text{ Ra}^{-0.314}$	$0.134 \text{ Ra}^{-0.242}$
Rigid-free	$0.247 \text{ Ra}^{0.277}$	$1.626 \text{ Ra}^{-0.317}$	$0.655 \text{ Ra}^{-0.328}$
Free-free	$0.253 \text{ Ra}^{0.306}$	$1.191 \text{ Ra}^{-0.289}$	$0.243 \text{ Ra}^{-0.259}$

is, approximately as $\text{Ra}^{-0.3}$. Table A1 also shows scalings for dimensionless topography and dimensionless gravity, which all scale roughly as $\text{Ra}^{-0.3}$ although there are some differences in detail. This scaling can be understood in broad terms from the expectation that the dynamic topography should be proportional to the boundary layer thickness (Parsons & Daly, 1983). Since the scaling used in Table A1 includes a ΔT_p factor, the scalings of the dimensional RMS gravity and topography in Table 2 go roughly as $\text{Ra}^{-0.6}$.

The behavior of the power spectral density with Rayleigh number is illustrated in Figure A2 for the free-rigid simulations. The principal effect of changing Rayleigh number is to change the amplitude of the power spectral density. This can be understood from the scalings of RMS gravity with Rayleigh number above, and the Parseval's theorem result

$$\Delta g_{\text{RMS}}^2 = \frac{1}{2\pi} \int \phi_{\text{PSD}}^{\Delta g}(k) k dk \quad (\text{A10})$$

which relates the square of the RMS value Δg_{RMS} to its power spectral density $\phi_{\text{PSD}}^{\Delta g}(k)$. Since RMS gravity in Table 2 scales roughly as $\text{Ra}^{-0.6}$, the power spectral density would be expected to scale as $\text{Ra}^{-1.2}$. This effect can be seen in Figure A2: Reducing the Rayleigh number by a factor of 10 leads to a little over an order of magnitude shift in the amplitude of the spectra. In addition to the change in amplitude, there are some more subtle changes in the spectra associated with changing the Rayleigh number. The lower Rayleigh number simulations appear to have relatively higher power at long wavelengths than at short wavelengths. This is as expected from the nature of the boundary layers, which are thicker for the lower Rayleigh number runs.

A5. The Relationship Between Temperature and Potential Temperature

To perform melting calculations, it is necessary to convert from dimensionless potential temperature back to real temperature. In this section we describe this conversion and justify the approximate form of energy




Figure A2. Power spectral density of gravity anomalies at the top of the convecting box as in Figure 8a, but showing the variation with Rayleigh number for the free-rigid simulations.

conservation that has been used in (6). In dimensional variables, conservation of energy can be written

$$\rho T \frac{DS}{Dt} = k \nabla^2 T + \Psi \quad (\text{A11})$$

where S is the specific entropy of a fluid parcel, k is the thermal conductivity (assumed constant), and Ψ is the viscous dissipation. The potential temperature θ can be defined in differential form as

$$dS = c_p d(\log \theta) \quad (\text{A12})$$

where c_p is the specific heat capacity at constant pressure (also assumed constant). The energy equation (A11) becomes

$$\frac{D\theta}{Dt} = \kappa \frac{\theta}{T} \nabla^2 T, \quad (\text{A13})$$

where $\kappa = k/(\rho c_p)$ is the thermal diffusivity, and viscous dissipation has been neglected. The use of (6) as a dimensionless governing equation for potential temperature is justified provided the approximation

$$\frac{1}{\theta} \nabla^2 \theta \approx \frac{1}{T} \nabla^2 T \quad (\text{A14})$$

is accurate and that viscous dissipation is sufficiently small to be neglected.

A5.1. No Melting

In the absence of melting, the differential of potential temperature can be related to those of temperature and pressure through the standard relationship

$$c_p \frac{d\theta}{\theta} = dS = c_p \frac{dT}{T} - \frac{\alpha}{\rho} dP \quad (\text{A15})$$

where α is the thermal expansivity. The principal variation in pressure is hydrostatic. Writing $dP = -\rho g dz$, (A15) can be written in terms of temperature and depth as

$$\frac{d\theta}{\theta} = \frac{dT}{T} + \frac{dz}{h_a} \quad (\text{A16})$$

where h_a is the adiabatic scale height, defined by $h_a = c_p/(\alpha g) \approx 3,300$ km. Integration of (A16) yields the relationship between temperature and potential temperature in regions which are not partially molten,

$$T = \theta \exp\left(\frac{z_{\text{ref}} - z}{h_a}\right) \quad (\text{A17})$$

where z_{ref} is a reference depth, the depth at which potential temperature is chosen to be equal to real temperature. This depth is chosen to be the Earth's surface in this work.

From (A16) it follows that

$$\frac{1}{\theta} \nabla^2 \theta = \frac{1}{T} \nabla^2 T + \frac{2}{h_a} \frac{1}{T} \frac{\partial T}{\partial z} + \frac{1}{h_a^2}. \quad (\text{A18})$$

The magnitude of the second and third terms on the right-hand side relative to the first term scales approximately as l/h_a and $(l/h_a)^2$ where l is a typical scale over which the temperature varies. If that scale l were the whole of the convecting layer, then $l/h_a = d/h_a = 0.18$ (a parameter known as the dissipation number), which is relatively small. In fact, the length scale of the vertical temperature variations will be much smaller than the layer depth, with boundary layer thicknesses on the order of 100 km or less, giving $l/h_a = 0.03$. Thus, both the second and third terms on the right-hand side of (A18) are sufficiently small that the approximation in (A14) is well justified in the regions that are not partially molten (McKenzie, 1970). The small dissipation number for upper mantle convection also justifies the neglect of viscous dissipation term in (A11).

An additional approximation has been made in writing the buoyancy term on the right-hand side of the Stokes equation in (2) in terms of the potential temperature θ . Formally, density variations in the fluid are determined by the actual temperature, not potential temperature, and the right-hand side of (2) should be $-\rho_0 g \alpha T \hat{z}$. The convective flow is driven by horizontal gradients in the actual temperature, not the potential

temperature. From (A17) it follows that the horizontal gradients in temperature are related to the horizontal gradients in potential temperature by

$$\frac{\partial T}{\partial x} = \frac{\partial \theta}{\partial x} \exp\left(\frac{z_{\text{ref}} - z}{h_a}\right), \quad (\text{A19})$$

with a similar expression for the y derivative. The horizontal gradients of potential temperature and actual temperature differ by an exponential factor whose magnitude is at most the exponential of the dissipation number. For the upper mantle convection we consider here, this is a relatively small difference and justifies the approximation made in using the potential temperature in (2).

A5.2. Melting

The convection simulations provide a 3-D grid of potential temperature (entropy) within the box. To turn this into melting rate, the hydrous melting parametrization of Katz et al. (2003) was used to calculate the expected degree of melting F at each grid point assuming isentropic decompression melting to the given potential temperature and pressure at each grid point. The original parametrization of degree of melting in Katz et al. (2003) is given with pressure and temperature as the thermodynamic variables. This parametrization can be recast in terms of pressure and entropy (or potential temperature) by numerically integrating the relevant differential expressions. The differential expression for entropy when melting is

$$c_p \frac{d\theta}{\theta} = dS = c_p \frac{dT}{T} - \left(\frac{\alpha}{\rho}\right) dP + \Delta S dF, \quad (\text{A20})$$

where F is the degree of melting, ΔS is the specific entropy difference between the two phases, and

$$\left(\frac{\alpha}{\rho}\right) = F \frac{\alpha_f}{\rho_f} + (1 - F) \frac{\alpha_s}{\rho_s}. \quad (\text{A21})$$

The Katz et al. (2003) parametrization accounts for the different thermal expansivities α_s, α_f ; and densities ρ_s, ρ_f of the two phases (solid and liquid respectively), but the specific heat c_p is assumed identical for both phases. All parameter values used here are identical to those in Table 2 of Katz et al. (2003), with the exception of the specific entropy difference between the two phases which we set as $\Delta S = 400 \text{ J kg}^{-1} \text{ K}^{-1}$. The parametrization provides the degree of melting F as a function of temperature T and pressure P which can be expressed in differential form as

$$dF = \left(\frac{\partial F}{\partial T}\right)_P dT + \left(\frac{\partial F}{\partial P}\right)_T dP. \quad (\text{A22})$$

Given a parcel of material that is subsolidus at a given potential temperature θ and depth z , (A17) gives the relationship between temperature and depth (or pressure) throughout the subsolidus region. Once the material crosses the solidus, the relationship between temperature and pressure at constant entropy can be obtained by numerically integrating equation (A20), (A21), and (A22) with $dS = 0$. Knowing the temperature and pressure then allows F to be calculated. From a series of constant entropy integrations for different potential temperatures a parametrization of F as a function of entropy and pressure was generated. Using this entropy parametrization, the grid of potential temperature and depth values in the box were converted to a grid of F values. This grid of F was then converted to a melting rate Γ using

$$\Gamma = \frac{\partial F}{\partial t} + \mathbf{v} \cdot \nabla F \quad (\text{A23})$$

where \mathbf{v} is the velocity. The time derivative was calculated using a first-order accurate finite difference approximation. The spatial gradient was calculated using a second-order accurate finite difference approximation. The melting rate Γ was vertically integrated to produce the plots shown in Figures 9 and 10. Only those regions where the melt rate was positive (i.e., melting) were included in the vertical integral. In the calculation of these figures, the advective term ($\mathbf{v} \cdot \nabla F$) was larger by more than an order of a magnitude than the time-dependent term ($\partial F / \partial t$), and an excellent approximation to the melting rates can be obtained from the advective term alone.

The calculation of melting rates was performed here as a postprocessing operation after running standard single-phase convection simulations. We assume that all the melt that is generated moves to the surface and

that none remains in the source region to freeze as the mantle material cools. It should be noted that this calculation neglects potentially important back effects that melt can have on the flow, for example, arising from the buoyancy of the melt, and the thermal effects of the consumption of latent heat. Indeed, in regions where melt is present, the approximation in (A14) can cease to be good approximation. However, the melting regions are only a small proportion of the overall domain, and the changes in temperature due to melting are small. For the free-rigid case with a lithospheric thickness of 80 km the average temperature change in the regions undergoing melting is 10 °C, and the maximum change in the whole box is 90 °C. Furthermore, such changes in temperature occur within the thermal boundary layer where heat is transported by conduction and where the temperature variations are large whether or not melting occurs. The effect of the temperature changes resulting from melting on the large-scale dynamics is therefore negligible.

There are other back effects of melt extraction on the convective flow that have been neglected. When melt is extracted from the mantle, the remaining residue has a different density than it had before the melt was extracted. There is thus the potential for this depletion by melting to change buoyancy forces and hence the flow. However, the density changes in the residue are small. Even for 20% melt extraction the relative density changes on depletion are on the order of -0.5% (Schutt & Leshner, 2006), equivalent to a density change from temperature variations of 125 °C.

Another potential back effect that has been neglected arises from the effect of volcanic loading on melt production rates. That changes in loading at the Earth's surface can influence melt production rates is well known from studies that have looked at the volcanic response to changes in ice cover in Iceland (Eksinichol et al., 2019; Jull & McKenzie, 1996; MacLennan et al., 2002). Beneath Iceland most melt generation occurs in the upper 100 km of the mantle, where the upwelling rate, of about 10 mm/yr, is driven by the separating plates. The thickness of the ice on Iceland reached about 3 km, equivalent to a thickness of rock of about 1 km, over about 10^5 years, corresponding to an equivalent rock accumulation rate of 10 mm/yr. Melt generation therefore ceases during the construction of the icecap, and all the melt that would normally have been generated during 10^5 years is instead released when the ice melts, in about 10^3 years. The behavior of melt generation within the convecting region beneath the lithosphere is very different. For the free-rigid case with a lithospheric thickness of 80 km shown in Figure 9, the upwelling rate of the solid mantle where the melt generation rate is fastest is 27 mm/yr. The accumulation rate of melt at the surface is about 300 m/Ma, or 0.3 mm/yr. This rate is therefore about 1/100 of the upwelling rate and will have no significant effect on the melt generation rate. Moreover, volcanic loads at the surface will be eroded over time, and their influence on the mantle beneath is further attenuated by the finite elastic strength of the overlying lithosphere.

Acknowledgments

The basis of this paper is Matthew Lees's Part III project in Earth Sciences. We thank Harro Schmeling and an anonymous reviewer for their comments which improved the paper. We thank Julianne Dannberg and Mark Hoggard for their advice. We thank the Leverhulme Trust for support. This work was performed using resources provided by the Cambridge Service for Data Driven Discovery (CSD3) operated by the University of Cambridge Research Computing Service (www.csd3.cam.ac.uk), provided by Dell EMC and Intel using Tier-2 funding from the Engineering and Physical Sciences Research Council (Capital Grant EP/P020259/1), and DiRAC funding from the Science and Technology Facilities Council (www.dirac.ac.uk). No new data have been generated in this work. The code for the numerical simulations of convection is available online (at aspect.geodynamics.org).

References

- Aubaud, C., Hauri, E. H., & Hirschmann, M. M. (2004). Hydrogen partition coefficients between nominally anhydrous minerals and basaltic melts. *Geophysical Research Letters*, 31, L20611. <https://doi.org/10.1029/2004GL021341>
- Ball, P. W., White, N. J., Masoud, A., Nixon, S., Hoggard, M. J., MacLennan, J., et al. (2019). Quantifying asthenospheric and lithospheric controls on mafic magmatism across North Africa. *Geochemistry, Geophysics, Geosystems*, 20, 3520–3555. <https://doi.org/10.1029/2019GC008303>
- Bangerth, W., Dannberg, J., Gassmüller, R., Heister, T., & Others (2018). *ASPECT v2.0.1 [software]*. Davis, CA: Computational Infrastructure for Geodynamics.
- Bruinsma, S. L., Förste, C., Abrikosov, O., Lemoine, J.-M., Marty, J.-C., Mulet, S., et al. (2014). ESA's satellite-only gravity field model via the direct approach based on all GOCE data. *Geophysical Research Letters*, 41, 7508–7514. <https://doi.org/10.1002/2014GL062045>
- Burke, K. (1996). The African plate. *South African Journal of Geology*, 99(4), 341–409.
- Busse, F. H., & Whitehead, J. A. (1971). Instabilities of convection rolls in a high Prandtl number fluid. *Journal of Fluid Mechanics*, 47(2), 305–320.
- Busse, F. H., & Whitehead, J. A. (1974). Oscillatory and collective instabilities in large Prandtl number convection. *Journal of Fluid Mechanics*, 66(1), 67–79.
- Christensen, U., & Harder, H. (1991). 3-D convection with variable viscosity. *Geophysical Journal International*, 104(1), 213–220.
- Craig, C. H., & McKenzie, D. (1987). Surface deformation, gravity and the geoid from a three-dimensional convection model at low Rayleigh numbers. *Earth and Planetary Science Letters*, 83(1–4), 123–136.
- Crosby, A. G., & McKenzie, D. (2009). An analysis of young ocean depth, gravity and global residual topography. *Geophysical Journal International*, 178(3), 1198–1219.
- Crosby, A. G., McKenzie, D., & Sclater, J. G. (2006). The relationship between depth, age and gravity in the oceans. *Geophysical Journal International*, 166(2), 553–573.
- Dannberg, J., & Heister, T. (2016). Compressible magma/mantle dynamics: 3-D, adaptive simulations in ASPECT. *Geophysical Journal International*, 207(3), 1343–1366. <https://doi.org/10.1093/gji/ggw329>
- Eason, D. E., Sinton, J. M., Grönvold, K., & Kurz, M. D. (2015). Effects of deglaciation on the petrology and eruptive history of the Western Volcanic Zone, Iceland. *Bulletin of Volcanology*, 77(6), 47.
- Ebinger, C. J., & Sleep, N. H. (1998). Cenozoic magmatism throughout east Africa resulting from impact of a single plume. *Nature*, 395(6704), 788–791.

- Eksinichol, I., Rudge, J. F., & MacLennan, J. (2019). Rate of melt ascent beneath Iceland from the magmatic response to deglaciation. *Geochemistry, Geophysics, Geosystems*, 20, 2585–2605. <https://doi.org/10.1029/2019GC008222>
- Erlank, A. J., Waters, F. G., Hawkesworth, C. J., Haggerty, S. E., Allsopp, H. L., Rickard, R. S., & Menzies, M. A. (1987). Evidence for mantle metasomatism in peridotite nodules from the Bultfontein Floors, Kimberley, South Africa. In M. A. Menzies, & C. J. Hawkesworth (Eds.), *Mantle metasomatism* (pp. 221–311). Academic Press, London.
- Fitton, J. G. (1987). The Cameroon line, West Africa: A comparison between oceanic and continental alkaline volcanism. *Geological Society, London, Special Publications*, 30(1), 273–291.
- Flament, N., Gurnis, M., & Müller, R. D. (2013). A review of observations and models of dynamic topography. *Lithosphere*, 5(2), 189–210.
- Frost, H. J., & Ashby, M. F. (1982). *Deformation-mechanism maps: The plasticity and creep of metals and ceramics*. Oxford: Pergamon Press.
- Galsa, A., & Lenkey, L. (2007). Quantitative investigation of physical properties of mantle plumes in three-dimensional numerical models. *Physics of Fluids*, 19(11), 116601.
- Hager, B. H., & Richards, M. A. (1989). Long-wavelength variations in Earth's geoid: Physical models and dynamical implications. *Philosophical Transactions of the Royal Society A*, 328(1599), 309–327.
- Hasterok, D. (2013). A heat flow based cooling model for tectonic plates. *Earth and Planetary Science Letters*, 361, 34–43.
- Heister, T., Dannberg, J., Gassmöller, R., & Bangerth, W. (2017). High accuracy mantle convection simulation through modern numerical methods—II: Realistic models and problems. *Geophysical Journal International*, 210(2), 833–851.
- Hoggard, M. J., White, N., & Al-Attar, D. (2016). Global dynamic topography observations reveal limited influence of large-scale mantle flow. *Nature Geoscience*, 9(6), 456–463.
- Hoggard, M. J., Winterbourne, J., Czarnota, K., & White, N. (2017). Oceanic residual depth measurements, the plate cooling model, and global dynamic topography. *Journal of Geophysical Research: Solid Earth*, 122, 2328–2372. <https://doi.org/10.1002/2016JB013457>
- Holmes, A. (1965). *Principles of physical geology*. London: Nelson.
- Houseman, G. A. (1990). The thermal structure of mantle plumes: Axisymmetric or triple-junction? *Geophysical Journal International*, 102(1), 15–24.
- Jarvis, G. T., & McKenzie, D. P. (1980). Convection in a compressible fluid with infinite Prandtl number. *Journal of Fluid Mechanics*, 96(3), 515–583.
- Jenkins, D. R. (1988). Interpretation of shadowgraph patterns in Rayleigh-Bénard convection. *Journal of Fluid Mechanics*, 190(-1), 451.
- Johnson, Kevin T. M., Dick, Henry J. B., & Shimizu, N. (1990). Melting in the oceanic upper mantle: An ion microprobe study of diopsides in abyssal peridotites. *Journal of Geophysical Research*, 95(B3), 2661.
- Jones, S. M., Lovell, B., & Crosby, A. G. (2012). Comparison of modern and geological observations of dynamic support from mantle convection. *Journal of the Geological Society, London*, 169(6), 745–758.
- Jull, M., & McKenzie, D. (1996). The effect of deglaciation on mantle melting beneath Iceland. *Journal of Geophysical Research*, 101(B10), 21,815–21,828.
- Katz, R. F., Spiegelman, M., & Langmuir, C. H. (2003). A new parameterization of hydrous mantle melting. *Geochemistry, Geophysics, Geosystems*, 4(9), 1073. <https://doi.org/10.1029/2002GC000433>
- King, S. D., & Anderson, D. L. (1998). Edge-driven convection. *Earth and Planetary Science Letters*, 160(3–4), 289–296.
- Kokfelt, T. F., Hoernle, K., & Hauff, F. (2003). Upwelling and melting of the Iceland plume from radial variation of ^{238}U – ^{230}Th disequilibria in postglacial volcanic rocks. *Earth and Planetary Science Letters*, 214(1–2), 167–186.
- Koornneef, J. M., Stracke, A., Bourdon, B., & Grönvold, K. (2012). The influence of source heterogeneity on the U-Th-Pa-Ra disequilibria in post-glacial tholeiites from Iceland. *Geochimica et Cosmochimica Acta*, 87, 243–266.
- Kvernfold, O. (1979). Rayleigh-Bénard convection with one free and one rigid boundary. *Geophysical & Astrophysical Fluid Dynamics*, 12(1), 273–294.
- Larsen, T. B., Yuen, D. A., Moser, J., & Fornberg, B. (1997). A high-order finite-difference method applied to large Rayleigh number mantle convection. *Geophysical & Astrophysical Fluid Dynamics*, 84(1–2), 53–83.
- Lee, D.-C., Halliday, A. N., Fitton, J. G., & Poli, G. (1994). Isotopic variations with distance and time in the volcanic islands of the Cameroon line: Evidence for a mantle plume origin. *Earth and Planetary Science Letters*, 123(1–3), 119–138.
- Liang, Y., & Peng, Q. (2010). Non-modal melting in an upwelling mantle column: Steady-state models with applications to REE depletion in abyssal peridotites and the dynamics of melt migration in the mantle. *Geochimica et Cosmochimica Acta*, 74(1), 321–339.
- MacLennan, J., Jull, M., McKenzie, D., Slater, L., & Grönvold, K. (2002). The link between volcanism and deglaciation in Iceland. *Geochemistry, Geophysics, Geosystems*, 3(11), 1062. <https://doi.org/10.1029/2001GC000282>
- McKenzie, D. (1970). Temperature and potential temperature beneath island arcs. *Tectonophysics*, 10(1–3), 357–366.
- McKenzie, D. (1985a). ^{230}Th – ^{238}U disequilibrium and the melting processes beneath ridge axes. *Earth and Planetary Science Letters*, 72(2–3), 149–157.
- McKenzie, D. (1985b). The extraction of magma from the crust and mantle. *Earth and Planetary Science Letters*, 74(1), 81–91.
- McKenzie, D. (1994). The relationship between topography and gravity on Earth and Venus. *Icarus*, 112, 55–88.
- McKenzie, D. (2016). A note on estimating Te from Bouguer coherence. *GEM - International Journal on Geomathematics*, 7(1), 103–116.
- McKenzie, D., & Fairhead, D. (1997). Estimates of the effective elastic thickness of the continental lithosphere from Bouguer and free air gravity anomalies. *Journal of Geophysical Research*, 102(B12), 27,523–27,552.
- McKenzie, D., Roberts, J. M., & Weiss, N. O. (1974). Convection in the Earth's mantle: Towards a numerical simulation. *Journal of Fluid Mechanics*, 62(3), 465.
- McKenzie, D., & Weiss, N. (1975). Speculations on the thermal and tectonic history of the Earth. *Geophysical Journal International*, 42(1), 131–174.
- McKenzie, D., Yi, W., & Rummel, R. (2014). Estimates of Te from GOCE data. *Earth and Planetary Science Letters*, 399, 116–127.
- McKenzie, D., Yi, W., & Rummel, R. (2015). Estimates of Te for continental regions using GOCE gravity. *Earth and Planetary Science Letters*, 428, 97–107.
- Michael, P. (1995). Regionally distinctive sources of depleted MORB: Evidence from trace elements and H₂O. *Earth and Planetary Science Letters*, 131(3–4), 301–320.
- Milelli, L., Fourel, L., & Jaupart, C. (2012). A lithospheric instability origin for the Cameroon Volcanic Line. *Earth and Planetary Science Letters*, 335–336, 80–87.
- Morgan, W. J. (1971). Convection plumes in the lower mantle. *Nature*, 230(5288), 42–43.
- Parsons, B., & Daly, S. (1983). The relationship between surface topography, gravity anomalies, and temperature structure of convection. *Journal of Geophysical Research*, 88(B2), 1129.
- Priestley, K., McKenzie, D., & Ho, T. (2018). A lithosphere-asthenosphere boundary—A global model derived from multimode surface-wave tomography and petrology, *Lithospheric discontinuities* (pp. 111–123). American Geophysical Union (AGU).

- Rexer, M., & Hirt, C. (2015). Spectral analysis of the Earth's topographic potential via 2D-DFT: A new data-based degree variance model to degree 90,000. *Journal of Geodesy*, 89(9), 887–909.
- Richter, F. M., & Parsons, B. (1975). On the interaction of two scales of convection in the mantle. *Journal of Geophysical Research*, 80(17), 2529–2541.
- Saal, A. E., Hauri, E. H., Langmuir, C. H., & Perfit, M. R. (2002). Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. *Nature*, 419(6906), 451–455.
- Schubert, G., Turcotte, D. L., & Olson, P. (2001). *Mantle convection in the Earth and planets*. Cambridge, New York, Melbourne: Cambridge University Press.
- Schutt, D. L., & Leshner, C. E. (2006). Effects of melt depletion on the density and seismic velocity of garnet and spinel lherzolite. *Journal of Geophysical Research*, 111, B05401. <https://doi.org/10.1029/2003JB002950>
- Segel, L. A., & Stuart, J. T. (1962). On the question of the preferred mode in cellular thermal convection. *Journal of Fluid Mechanics*, 13(2), 289.
- Slater, L., McKenzie, D., Grönvold, K., & Shimizu, N. (2001). Melt generation and movement beneath Theistareykir, NE Iceland. *Journal of Petroleum*, 42(2), 321–354.
- Sleep, N. H. (1997). Lateral flow and ponding of starting plume material. *Journal of Geophysical Research*, 102(B5), 10,001–10,012.
- Sleep, N. H. (2008). Channeling at the base of the lithosphere during the lateral flow of plume material beneath flow line hot spots. *Geochemistry, Geophysics, Geosystems*, 9, Q08005. <https://doi.org/10.1029/2008GC002090>
- Sotin, C., & Labrosse, S. (1999). Three-dimensional thermal convection in an iso-viscous, infinite Prandtl number fluid heated from within and from below: Applications to the transfer of heat through planetary mantles. *Physics of the Earth and Planetary Interiors*, 112(3–4), 171–190.
- Spiegel, E. A., & Veronis, G. (1960). On the Boussinesq approximation for a compressible fluid. *The Astrophysical Journal*, 131, 442.
- Stracke, A., Bourdon, B., & McKenzie, D. (2006). Melt extraction in the Earth's mantle: Constraints from U-Th-Pa-Ra studies in oceanic basalts. *Earth and Planetary Science Letters*, 244(1–2), 97–112.
- Suh, C. E., Ayonghe, S. N., Sparks, R. S. J., Annen, C., Fitton, J. G., Nana, R., & Luckman, A. (2003). The 1999 and 2000 eruptions of Mount Cameroon: Eruption behaviour and petrochemistry of lava. *Bulletin of Volcanology*, 65(4), 267–281.
- Tackley, P. J. (1993). Effects of strongly temperature-dependent viscosity on time-dependent, three-dimensional models of mantle convection. *Geophysical Research Letters*, 20(20), 2187–2190.
- Tackley, P. J. (1996). Effects of strongly variable viscosity on three-dimensional compressible convection in planetary mantles. *Journal of Geophysical Research*, 101(B2), 3311–3332.
- Thorpe, R. S., & Smith, K. (1974). Distribution of Cenozoic volcanism in Africa. *Earth and Planetary Science Letters*, 22(1), 91–95.
- Travis, B., Weinstein, S., & Olson, P. (1990). Three-dimensional convection planforms with internal heat generation. *Geophysical Research Letters*, 17(3), 243–246.
- Turner, S., Kokfelt, T., Hoernle, K., Lundstrom, C., & Hauff, F. (2016). ²³¹Pa systematics in postglacial volcanic rocks from Iceland. *Geochimica et Cosmochimica Acta*, 185, 129–140.
- Vilella, K., & Deschamps, F. (2017). Thermal convection as a possible mechanism for the origin of polygonal structures on Pluto's surface. *Journal of Geophysical Research: Planets*, 122, 1056–1076. <https://doi.org/10.1002/2016JE005215>
- Vilella, K., & Deschamps, F. (2018). Temperature and heat flux scaling laws for isoviscous, infinite Prandtl number mixed heating convection. *Geophysical Journal International*, 214(1), 265–281.
- Warren, J. M. (2016). Global variations in abyssal peridotite compositions. *Lithos*, 248–251, 193–219.
- Weinstein, S. A., & Christensen, U. (1991). Convection planforms in a fluid with a temperature-dependent viscosity beneath a stress-free upper boundary. *Geophysical Research Letters*, 18(11), 2035–2038.
- White, D. B. (1988). The planforms and onset of convection with a temperature-dependent viscosity. *Journal of Fluid Mechanics*, 191(-1), 247.
- Whitehead, J. A., & Parsons, B. (1977). Observations of convection at Rayleigh numbers up to 760,000 in a fluid with large Prandtl number. *Geophysical & Astrophysical Fluid Dynamics*, 9(1), 201–217.
- Wilson, J. T. (1963). A possible origin of the Hawaiian Islands. *Canadian Journal of Physics*, 41(6), 863–870.
- Zhong, S. (2005). Dynamics of thermal plumes in three-dimensional isoviscous thermal convection. *Geophysical Journal International*, 162(1), 289–300.